

Supplementary table 1: Different genotypes of Apo B VNTR based on five-allelic model

S. No	Genotype	Patients (n=230)	Controls (n=300)	p value
1	1/1	1(0.4%)	2(0.6%)	ns
2	1/2	6(2.6%)	13(4.3%)	ns
3	1/3	3(1.3%)	16(5.3%)	ns
4	1/4	3(1.3%)	3(1.0%)	ns
5	1/5	22(9.6%)	19(6.3%)	ns
6	2/2	11(4.8%)	26(8.6%)	ns
7	2/3	8(3.5%)	48(16.0%)	0.0001
8	2/4	6(2.6%)	12(4.0%)	ns
9	2/5	19(8.3%)	6(2.0%)	0.0001
10	3/3	44(19.1%)	64(21.3%)	ns
11	3/4	19(8.3%)	18(6.0%)	ns
12	3/5	31(13.5%)	35(11.6%)	ns
13	4/4	25(10.9%)	18(6.0%)	ns
14	4/5	12(5.2%)	9(3.0%)	ns
15	5/5	20(8.7 %)	11(3.6 %)	0.02

Supplementary Figure 1: Different alleles at ApoB 3' HVR with sizing ladder.

Lane 1: Allelic ladder for ApoB3' HVR, Lane 2–6: different samples showing various alleles of ApoB 3' HVR and lane 7: 50 bp ladder.

