

Optical coherence tomography on the right side: pulmonary vascular disease in mitral stenosis

A 59-year-old woman presented with a 6-month history of worsening fatigue and exertional dyspnoea. On physical examination, she had a mitral stenosis murmur, a loud P2, an irregular pulse and mild peripheral oedema. The electrocardiogram showed atrial fibrillation and the echocardiogram was notable for a fibro-calcified mitral valve, with severe restriction of leaflet mobility and an area of 1 cm². Biventricular function was normal. As the valve anatomy was considered suitable for percutaneous intervention, the patient was scheduled for a balloon valvuloplasty. The preintervention cardiac catheterisation showed a pulmonary artery pressure of 41/20/30 mm Hg. After successful mitral dilatation, optical coherence tomography (OCT) (LightLab Imaging Inc, Westford, Massachusetts, USA) was performed on the distal pulmonary artery (figure 1). It showed thickened vessel wall either in small vessels (<1 mm) (figure 2A) or larger vessels (diameter ca. 2 mm) (figure 2B). OCT is a potentially useful tool for unravelling the mechanisms of group 2 pulmonary hypertension.¹




Figure 1 Pulmonary angiogram showing the location of the optical coherence tomography probe on the right distal pulmonary artery. A Judkins right guiding catheter was used (Launcher, Medtronic, Minnesota, USA) for advancing the probe and contrast as flushing solution.

Elisabete Jorge,^{1,2} Henrique Faria,¹ Cristina Silva,¹ Manuel Pan³

¹Serviço de Cardiologia, Hospitais da Universidade de Coimbra, Centro Hospitalar e Universitário de Coimbra, Coimbra, Portugal

²Faculdade de Medicina, Universidade de Coimbra, Coimbra, Portugal

³Servicio de Cardiología, Hospital Reina Sofia, Córdoba, Spain




Figure 2 (A) At the level of the distal pulmonary artery (<1 mm² area), the vessel wall is markedly thickened (0.23 mm). (B) In this 2-mm wide distal pulmonary artery, the walls are also markedly thickened.

Correspondence to Dr Elisabete Sofia Azenha Balhau Jorge, Serviço de Cardiologia, Hospitais da Universidade de Coimbra, Centro Hospitalar e Universitário de Coimbra, Praceta Mota Pinto, Coimbra 3000-075, Portugal; elisabetejorge@gmail.com

Contributors EJ designed the project, performed the procedure and wrote the manuscript. HF performed the procedure and revised the draft manuscript. CS prepared and assisted the procedure. MP designed the project and revised the draft manuscript.

Competing interests None.

Patient consent Obtained.

Ethics approval Comissão de Ética da Faculdade de Medicina da Universidade de Coimbra.

Provenance and peer review Not commissioned; internally peer reviewed.

To cite Jorge E, Faria H, Silva C, *et al.* *Heart Asia* Published Online First: [please include Day Month Year] doi:10.1136/heartasia-2013-010321

Heart Asia 2013;0:85. doi:10.1136/heartasia-2013-010321

REFERENCE

- Li N, Zhang S, Hou J, *et al.* Assessment of pulmonary artery morphology by optical coherence tomography. *Heart Lung Circ* 2012;21:778–81.