East meets West: the influence of racial, ethnic and cultural risk factors on cardiac surgical risk model performance

Sarah Soo-Hoo, Samantha Nemeth, Onur Baser, Michael Argenziano, Paul Kurlansky

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/heartasia-2017-010995>).

Department of Surgery, Columbia University, New York, New York City, USA

Correspondence to

Dr Paul Kurlansky, Department of Surgery, Columbia University Medical Center, New York NY 10032, USA; pk2245@cumc.columbia.edu

Presented at the 2016 Scientific Sessions of the American Heart Association, New Orleans, 14 November 2016.

Received 27 December 2017
Revised 5 February 2018
Accepted 6 February 2018

ABSTRACT

Objective To explore the impact of racial and ethnic diversity on the performance of cardiac surgical risk models, the Chinese SinoSCORE was compared with the Society of Thoracic Surgeons (STS) risk model in a diverse American population.

Methods The SinoSCORE risk model was applied to 13 969 consecutive coronary artery bypass surgery patients from twelve American institutions. SinoSCORE risk factors were entered into a logistic regression to create a 'derived' SinoSCORE whose performance was compared with that of the STS risk model.

Results Observed mortality was 1.51% (66% of that predicted by STS model). The SinoSCORE 'low-risk' group had a mortality of $0.15\% \pm 0.04\%$, while the medium-risk and high-risk groups had mortalities of $0.35\% \pm 0.06\%$ and $2.13\% \pm 0.14\%$, respectively. The derived SinoSCORE model had a relatively good discrimination (area under of the curve (AUC)=0.785) compared with that of the STS risk score (AUC=0.811; $P=0.18$ comparing the two). However, specific factors that were significant in the original SinoSCORE but that lacked significance in our derived model included body mass index, preoperative atrial fibrillation and chronic obstructive pulmonary disease.

Conclusion SinoSCORE demonstrated limited discrimination when applied to an American population. The derived SinoSCORE had a discrimination comparable with that of the STS, suggesting underlying similarities of physiological substrate undergoing surgery. However, differential influence of various risk factors suggests that there may be varying degrees of importance and interactions between risk factors. Clinicians should exercise caution when applying risk models across varying populations due to potential differences that racial, ethnic and geographic factors may play in cardiac disease and surgical outcomes.

INTRODUCTION

Despite considerable improvements in cardiovascular therapies, coronary artery disease (CAD) remains the leading cause of morbidity and mortality worldwide.^{1–3} Likewise, this trend shows no signs of reversing in rapidly developing countries such as China, where there is expected to be a >50% increase in coronary events over the next 15 years.² This has resulted in a similar upwards shift in coronary artery bypass grafting (CABG) surgery volume. However, notably absent from this movement is the adoption of a standard risk model to assess CABG mortality in Chinese populations.¹⁴

Cardiac risk models have traditionally been derived from and developed for American and

European populations; however, these models may prove inaccurate when applied to Chinese populations.^{5–8} As medicine becomes increasingly collaborative and standardised internationally, it remains unclear if objective standards can be applied in risk adjustment to compare outcomes and prognosis in the setting of racial, ethnic and cultural differences. We therefore undertook the novel exploration of applying SinoSCORE, a contemporary Chinese cardiac risk model, in a diverse American population.^{9 10} Given both the increasing emphasis on quality assessment in cardiac surgery and the increasing prevalence of CAD in China, we find this investigation to be both timely and compelling.¹¹

MATERIALS AND METHODS

Study population

The study population consisted of all patients who underwent CABG surgery in any of 12 cardiac surgery programmes with the Columbia University HeartSource programme from May 2006 to March 2017. These 12 programmes represent a broad array of medical centres—academic and non-academic, urban, suburban and rural located throughout different regions in the USA. The purpose of the Columbia HeartSource project is to help community hospitals enhance an existing cardiovascular programme or to launch one de novo. The programme involves involvement of Columbia faculty for real-time consultation, discussion of challenging cases and external peer review of all deaths and other significant cases. As a result, Columbia HeartSource personnel have readily available access to affiliate's surgical data. All surgical cases are prospectively entered into a web-based proprietary Society of Thoracic Surgeons (STS)-compliant database that is regularly monitored for timeliness and accuracy.

Our hypothesis is that racial and ethnic differences between populations limit the ability of professionally accepted risk models to predict mortality following CABG surgery. More specifically, we hypothesise that differences between American and Chinese populations will impact the performance of risk models derived from each of those surgical populations. Ideally, our hypothesis would be tested by applying both a Chinese and an American risk model to a representative sample of patients from each population; however, since the Chinese population was not available for study, we elected to compare the performance of each risk model in a diverse American population.

To cite: Soo-Hoo S, Nemeth S, Baser O, et al. *Heart Asia* 2018;**10**:1–6. doi:10.1136/heartasia-2017-010995

A total of 13 969 consecutive patients underwent CABG surgery and could have an STS operative mortality risk score calculated (ie, isolated CABG or CABG+valve) during this study period. Preoperative and demographic information, operative data and in-hospital mortality for all patients were retrieved from the STS-compliant HeartSource database. Excluded were all patients who had associated procedures for which there was no STS risk prediction model.

Primary predictors

All risk factors relevant to SinoSCORE^{9 10} and STS^{12 13} risk calculations were collected. Available data were entered into each risk calculation using appropriate definitions. In cases where there were minor discrepancies between the definitions of various risk factors (chronic obstructive pulmonary disease (COPD), atrial fibrillation and so on) between the two models, reasonable approximations were made (see online supplementary file.) Based on original SinoSCORE methodology, a SinoSCORE was then calculated for each patient. STS mortality risk scores were retrieved from the HeartSource database.⁹ SinoSCORE was calculated for all patients undergoing CABG, including those with associated valve surgery, whereas separate STS risk models were employed for each isolated CABG and combined CABG/valve procedure.^{12 13} In accordance with original SinoSCORE methodology, patients were stratified into low-risk, medium-risk and high-risk groups according to their respective SinoSCOREs (low: <1, medium: 1 < and <6, high: ≥6).

Study endpoints

The primary endpoint for this study was mortality. All calculations for mortality related to SinoSCORE were based on in-hospital mortality—or death occurring during the hospitalisation in which the operation was performed—as SinoSCORE is a risk prediction model for in-hospital mortality. All calculations for mortality related to the STS risk models were based on operative mortality—or death occurring during the hospitalisation in which the operation was performed, even if after 30 days (including patients transferred to other acute care facilities), or death occurring after discharge from the hospital, but before the end of the thirtieth postoperative day—as STS is a risk prediction model for operative mortality.

Statistical analyses

Categorical data are presented as frequency distributions and simple percentages. Values of continuous variables are expressed as a mean ± SD or as a median ± IQR, as appropriate. Performance of both the SinoSCORE and STS risk models were assessed by comparing the observed and expected mortalities for the overall population and each subgroup, defined by their predicted risk scores.

To assess discrimination within the overall cohort and subgroups, we used receiver operator characteristics to calculate the area under the curve (AUC). An AUC of 0.5 is indicative of no discriminatory ability, while an AUC of 1.0 is considered to be perfect discriminatory ability. To assess calibration, we used the Hosmer-Lemeshow (H-L) goodness-of-fit test. H-L P values greater than 0.05 were considered to be well-calibrated models for our study population. Calibration plots comparing observed to expected mortality based on the predicted risks calculated by the SinoSCORE and STS were created. A perfectly calibrated prediction model would follow the 45° line, while curves below the line are considered to be overestimates and curves above the line are considered to be underestimates. All analyses were performed using SAS, V.9.2. P < 0.05 was considered significant.

The published form of the SinoSCORE risk model does not provide a direct correlation between SinoSCORE and predicted risk of mortality for each individual score, but rather categorises patients into low-risk, medium-risk and high-risk groups, indicating a range of predicted mortality for each of these subgroups. Due to the inability to assign a distinct predicted risk of mortality to each increment of the SinoSCORE, we developed a logistic regression model using risk factors found to be significant in the originally developed SinoSCORE as independent variables and hospital mortality as the dependent variable, resulting in a ‘derived SinoSCORE’. We then compared the performance of this ‘derived SinoSCORE’ with the STS risk model in our diverse American patient population.

Definition reconciliation

Data regarding COPD for the SinoSCORE risk model was replaced using single imputation methods. SinoSCORE defines COPD as ‘long-term use of bronchodilators or steroids for lung disease’; however, use of bronchodilators was not collected in earlier versions of the STS Database (V.2.52 and V.2.61) but was recorded in later versions (V.2.73 and V.2.81). Patients with applicable COPD information available were examined. For populations with no information available, an imputed value for SinoSCORE proportionate to the probability of the prevalence of COPD in that population was calculated and assigned to all patients with missing data.

Institutional review board

This study was approved by the Columbia University Institutional Review Board with waiver of consent.

RESULTS

Demographics

A summary of population characteristics can be found in table 1. The mean age of our patient cohort was 67.14, and 74.55% were men. Of note, only 1.38% of the study population was of Asian origin. The SinoSCORE ‘low risk’ group (≤1) had a mean SinoSCORE of -0.96 ± 1.22 , while the ‘medium’ ($1 < x < 6$) and ‘high’ (≥6) risk groups had a mean SinoSCORE of 3.4 ± 1.07 and 11.69 ± 4.47 , respectively.

Comparison in different populations (primary analyses)

Of the entire cohort, there were 211 deaths observed, resulting in an overall mortality of 1.51%. The STS observed to expected ratio for operative mortality was 0.66. The low-risk subgroup had a mortality of $0.15\% \pm 0.04\%$, the medium-risk subgroup had a mortality of $0.35\% \pm 0.06\%$ and the high-risk subgroup had a mortality $2.13\% \pm 0.14\%$. Observed and predictive operative mortality using SinoSCORE methodology in the American and Chinese populations are summarised in table 2. In both populations, SinoSCORE successfully identified patients that were low-risk, medium-risk and high-risk for CABG-related mortality. However, in contrast to the distinct non-overlapping CIs in the original Chinese population from which the SinoSCORE was derived, there was considerable overlap in CIs between risk groups in the American population, suggesting that the discrimination of SinoSCORE in the American population may not be optimal.

Comparison of models (secondary analyses)

To explore whether this discrepancy in performance was a byproduct of the risk factors included in the SinoSCORE model or the population from which the model is derived,

Table 1 Baseline SinoSCORE and STS characteristics

SinoSCORE characteristics	n=13 969 (%)
Male	10 414 (74.55)
Age (years)	
<65	5319 (38.08)
65–69	2613 (18.71)
70–74	2324 (16.64)
≥75	3713 (26.58)
BMI	
≥24	11 294 (80.85)
<18	60 (0.43)
LVEF <50%	4342 (31.08)
NYHA	
Stage III	1567 (11.22)
Stage IV	786 (5.63)
Chronic renal failure	377 (2.70)
Extracardiac arteriopathy	3555 (25.45)
COPD	1228 (8.79)
Preoperative atrial fibrillation or flutter	1213 (8.68)
Preoperative critical stage	1226 (8.78)
Non-elective surgery	7631 (54.63)
Combined valve surgery	2233 (15.99)
STS characteristics	n=13 969 (%)
Age (years)	
<65	5319 (38.08)
65–69	2613 (18.71)
70–74	2324 (16.64)
≥75	3713 (26.58)
Diabetes	5723 (40.97)
Control: diet	883 (6.32)
Control: other	2879 (20.61)
Control: insulin	1961 (14.04)
Chronic renal failure	377 (2.70)
Chronic lung disease	1385 (9.91)
Immunocompromise	502 (3.59)
Peripheral arterial disease	1983 (14.20)
Cerebrovascular disease	2231 (15.97)
Prior CVA	882 (6.31)
Prior CABG	351 (2.51)
Prior valve	72 (0.52%)
Previous PCI	3809 (27.27)
MI when ≤24 hours	480 (3.44)
MI when 1–7 days	2757 (19.74)
MI when 7+ days	3099 (22.18)
CHF	3162 (22.64)
NYHA class	
No NYHA	10 102 (72.32)
NYHA Stage I	429 (3.07)
NYHA Stage II	1077 (7.71)
NYHA Stage III	1567 (11.22)
NYHA Stage IV	786 (5.63)
Cardiac symptoms at admission	
No symptoms or angina	1883 (14.35)
Symptoms unlikely to be ischaemia	381 (2.90)
Angina equivalent	285 (2.17)
Stable angina	2739 (20.88)
Unstable angina	4420 (33.69)
NSTEMI	2606 (19.86)

Continued

Table 1 Continued

STS characteristics	n=13 969 (%)
STEMI	530 (4.04)
Other	250 (1.91)
Cardiac symptoms at surgery	4774 (17)
No symptoms or angina	831 (17.41)
Angina equivalent	278 (5.82)
Stable angina	555 (11.63)
Unstable angina	1831 (38.35)
NSTEMI	881 (18.45)
STEMI	149 (3.12)
Other	249 (5.22)
Cardiac shock	254 (1.82)
Resuscitation yes	90 (0.64)
Atrial fibrillation	1213 (8.68)
Inotropes yes	182 (1.30)
Number of diseased vessels less than 3	3713 (26.58)
Number of diseased vessels 3	10 235 (73.27)
Ejection fraction	4342 (31.08)
Endocarditis type active	25 (0.18)
Endocarditis type treated	31 (0.22)
Mitral insufficiency	1293 (9.26)
Mitral stenosis	155 (1.11)
Tricuspid insufficiency	435 (3.11)
Incidence: first surgery	13 556 (97.04)
Incidence: reoperation	413 (2.96)
Status	
Elective	6337 (45.36)
Urgent	7095 (50.79)
Emergent	522 (3.74)
Salvage	14 (0.10)
IABP	
Preoperative	910 (6.51)
Intraoperative	269 (1.93)
Postoperative	52 (0.37)
Operative death	259 (1.85)

BMI, body mass index; CABG, coronary artery bypass grafting; CHF, congestive heart failure; COPD, chronic obstructive pulmonary disease; CVA, cerebrovascular accident; IABP, intra-aortic balloon pump; LVEF, left ventricular ejection fraction; MI, myocardial infarction; NSTEMI, non-ST elevation myocardial infarction; NYHA, New York Heart Association; PCI, percutaneous coronary intervention; STEMI, ST elevation myocardial infarction; STS, Society of Thoracic Surgeons.

we developed a model ('derived SinoSCORE') using logistic regression. Significant risk factors included in both the SinoSCORE and derived SinoSCORE model are summarised in [table 3](#).¹⁰ Factors that were found to be significant in the original SinoSCORE, but lacked significance in the derived SinoSCORE model included body mass index, preoperative atrial fibrillation, COPD and non-elective surgery.

The derived SinoSCORE model had a relatively good discrimination (AUC=0.785) for our patient cohort compared with that of the STS risk score (AUC=0.811), with no difference in discrimination between these two models (P=0.18; [figure 1](#)). Model calibration was evaluated using the H-L test. The derived SinoSCORE was well calibrated for this population (P=0.146), whereas the STS Risk Score was tended to overestimate operative mortality (P<0.001) due to the lower than expected mortality in this high-quality surgical population ([figure 2](#)).

Table 2 Mortality stratified by SinoSCORE risk categories: comparison of American study population with original Chinese SinoSCORE population

SinoSCORE	SinoSCORE American population			CIs (expected)	Original SinoSCORE Chinese population	
	N	Mean observed	SD		Observed	Expected
≤1 (low risk)	1341	0.15% (2/1341)	0.04	0.0% to 0.36%	0.37–0.81	0.52–0.89
2–5 (medium risk)	3394	0.35% (12/3394)	0.06	0.15% to 0.55%	1.13–1.98	1.29–1.94
≥6 (high risk)	9234	2.13% (197/9234)	0.14	1.84% to 2.43%	5.00–7.89	6.36–6.63

DISCUSSION

To our knowledge, this is the first study that applies the SinoSCORE to a non-Chinese population. With increasing globalisation of information and medical knowledge, these findings sound a certain note of caution in applying risk models to populations different from those from which they were derived. A stronger understanding of the underlying factors that limit the broad application of risk models may shed insight into the intrinsic differences between populations and reduce inappropriate utilisation of risk models, thereby potentially reducing patient mortality.^{8 14 15} Moreover, since China in particular has recently been more open to embracing Western medical standards, a more nuanced understanding of these limitations is increasingly relevant.

Application of the SinoSCORE model to the American population suggested questionable discrimination—as demonstrated by overlapping confidence intervals—and thus may not be an appropriate risk model for American populations. On further analysis, when SinoSCORE significant risk factors were applied to our cohort, the risk model (derived SinoSCORE) had a discrimination that was comparable with that of the STS ($P=0.18$). This suggests that there are similarities in the patient-related factors that determine operative risk, but that the relative influence of each risk factor may vary between populations (table 3).

Table 3 ORs of significant risk factors in SinoSCORE and derived SinoSCORE

Variable	SinoSCORE OR (95% CI) ¹¹	Derived SinoSCORE* OR (95% CI)
Age (years)	1.06 (1.04 to 1.08)	
65–69		1.69 (1.05 to 2.70)
70–74		1.72 (1.07 to 2.76)
≥75		2.95 (2.00 to 4.35)
BMI	0.91 (0.87 to 0.96)	
≥24		2.19 (0.64 to 8.49)
<18		0.78 (0.56 to 1.07)
LVEF <50%	0.97 (0.95 to 0.98)	1.85 (1.37 to 2.50)
NYHA	1.92 (1.43 to 2.58)	
Stage III		1.58 (1.09 to 2.28)
Stage IV		2.24 (1.50 to 3.35)
Chronic renal failure	2.4 (1.27 to 4.53)	1.62 (0.89 to 2.95)
Extracardiac arteriopathy	4.38 (2.30 to 8.33)	1.54 (1.16 to 2.05)
COPD	2.46 (1.36 to 4.45)	0.86 (0.63 to 1.16)
Preoperative atrial fibrillation or flutter	1.97 (1.33 to 2.92)	1.13 (2.07 to 4.14)
Preoperative critical stage	2.20 (1.29 to 3.75)	2.92 (2.07 to 4.14)
Non-elective surgery	2.38 (1.30 to 4.35)	1.20 (0.87 to 1.65)
Combined valve surgery	2.51 (1.76 to 3.57)	2.49 (1.83 to 3.40)

Bold font indicates $P<0.05$.

*Derived SinoSCORE was based on our population.

BMI, body mass index; COPD, chronic obstructive pulmonary disease; LVEF, left ventricular ejection fraction; NYHA, New York Heart Association.

Overlapping risk factors between SinoSCORE and derived SinoSCORE also suggest that risk factors for CABG mortality in both populations may be similar, but the differing ORs for identical risk factors indicate that the interaction between risk factors may be distinct between both populations. For example, chronic renal failure (OR: 2.4 vs 1.62), COPD (OR: 2.46 vs 0.86) and extracardiac arteriopathy (OR: 4.38 vs 1.54) played greater roles in mortality in the Chinese population than in the American population. This may be explained by underlying factors that have uniquely shaped the CAD landscape in China. For example, the prevalence of hypertension in China rose dramatically in the previous decade, and ineffective management of hypertension has been linked to the rising incidence of renal failure.^{16 17} The increased role of COPD in CABG mortality may reflect a higher incidence cigarette smoking and worsening air pollution, as pollutant concentrations have been recorded to be greater than four times the WHO recommended limit and has been linked to more than 1.3 million premature deaths in 2010.^{18–20} Moreover, China's urbanisation and industrialisation have been associated with the adoption of 'Western' dietary patterns and decreased physical activity that are likely linked to increasing rates of extracardiac arteriopathy.^{1 3 11 17 20} However, factors such as combined valve surgery (OR: 2.51 vs 2.49) and preoperative critical staging (OR: 2.20 vs 2.92) had similar impacts in both populations, which is not surprising given that these parameters are less influenced by cultural factors but rather tend to reflect sicker patients and more complicated operations. Our data shed light on the intrinsic differences between the Chinese and American populations included in this study, which is then reflected in the performance of risk models/risk scores. This study highlights how risk models may not capture underlying differences in non-clinical aspects of disease and supports the hypothesis that variation in patient composition may be a significant source of error in cardiac risk models.²¹

Figure 1 Receiver operator curves (ROCs) comparing 'derived' SinoSCORE with STS risk score models. STS, Society of Thoracic Surgeons.

Figure 2 Calibration plots comparing performance of 'derived' SinoSCORE with STS risk score models. STS, Society of Thoracic Surgeons.

While the derived SinoSCORE and STS had comparable discriminations, the derived SinoSCORE demonstrated a superior calibration to STS ($P=0.146$ vs $P<0.001$) in the setting of our patient population. This is unsurprising given the derived SinoSCORE was derived from the patient population to which it was applied and thus would understandably result in a superior calibration. In addition, the relatively high-performance network from which the data were derived from had an observed to expected mortality of 0.66 and likely contributed to STS model's overprediction of mortality and relatively poor calibration.

Our study highlights how risk model performance may be dependent on sample population and shaped by both clinical and non-clinical factors including racial and cultural differences,

which need to inform comparative performance assessment. Independent external validations of CABG risk models in broad populations should continue as current risk models may not be suitable for all populations.

Limitations

As alluded to in the Methods, the patient population would ideally have included both a Chinese and American population to evaluate risk model performance; however, we were unable to obtain to access this population. In addition, our cohort cannot definitively be said to represent the general American population. However, our sample included patients from a broad range of institutions that varied in size, location and volume. Likewise, SinoSCORE was derived from 42 centres that likely represent an appropriate sampling of the Chinese population with advanced CAD who undergo CABG. Furthermore, while definitions and criteria for certain risk factors did not align exactly between SinoSCORE and STS, appropriate approximations were made (see online supplementary file).

CONCLUSION

To our knowledge, this is the first study that applies the SinoSCORE to a non-Chinese population. The original SinoSCORE demonstrated limited discrimination when applied to an American population, indicating that it may not be appropriate for evaluating CABG mortality in certain populations. The derived SinoSCORE had a discrimination comparable with that of the STS, suggesting that there may be varying degrees of importance and interactions between risk factors within the context of the fundamental physiological similarity of patients undergoing advanced surgical coronary revascularisation. Clinicians must exercise caution when applying risk models across varying populations due to potential differences that racial, ethnic and geographic factors may play in cardiac disease and outcomes.

Contributors All authors participated in the analysis of the data and either manuscript writing or review.

Funding Funding for this project was supplied by entirely unrestricted internal funds from the Department of Surgery of Columbia University.

Disclaimer None of the authors has any conflict to disclose regarding any aspect of this work; all views expressed are those of the authors, not an official position of the Department of Surgery nor of the University.

Competing interests None declared.

Patient consent Not required.

Ethics approval Columbia University Human Subjects Protocol AAAQ2103.

Provenance and peer review Not commissioned; internally peer reviewed.

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- Chen WW, Gao RL, Liu LS, et al. China cardiovascular diseases report 2015: a summary. *J Geriatr Cardiol* 2017;14:1–10.
- Moran A, Gu D, Zhao D, et al. Future cardiovascular disease in China: markov model and risk factor scenario projections from the coronary heart disease policy model-china. *Circ Cardiovasc Qual Outcomes* 2010;3:243–52.
- Sanchis-Gomar F, Perez-Quilis C, Leischik R, et al. Epidemiology of coronary heart disease and acute coronary syndrome. *Ann Transl Med* 2016;4:256.
- Zheng Z, Li Y, Zhang S, et al. The Chinese coronary artery bypass grafting registry study: how well does the EuroSCORE predict operative risk for Chinese population? *Eur J Cardiothorac Surg* 2009;35:54–8.
- Bai Y, Wang L, Guo Z, et al. Performance of EuroSCORE II and SinoSCORE in Chinese patients undergoing coronary artery bypass grafting. *Interact Cardiovasc Thorac Surg* 2016;23:733–9.

Key messages

What is already known about this subject?

- To date, approximately 20 risk models have been developed to study mortality following coronary artery bypass surgery. Given the wealth of supporting data and penetration of the database in the American population, the Society of Thoracic Surgeons risk model has gained increasing stature as the most widely accepted standard in the USA. However, emergence of risk models from other countries has raised the question as to the potential limitations of applying risk models that were well derived in one population to other populations with differences in racial and ethnic composition.

What does this study add?

- Although application and limitations of American risk models for cardiac surgical mortality to Chinese patient populations has previously been reported, this is the first report of a Chinese model being applied to a diverse contemporary American population. The findings suggest that although there is a fundamental physiological similarity among patients undergoing coronary artery bypass surgery, the performance of risk models is limited by differences in the impact of various risk factors in different populations.

How might this impact on clinical practice?

- Although there is a tendency to seek 'international' standards for evaluating surgical performance, such efforts must always be tempered by adjustment for specific racial, ethnic and social differences between populations.

- 6 Sun C, Xu F, Liu X, *et al.* Comparison of validation and application on various cardiovascular disease mortality risk prediction models in Chinese rural population. *Sci Rep* 2017;7:43227.
- 7 Zheng X, Curtis JP, Hu S, *et al.* Coronary catheterization and percutaneous coronary intervention in China: 10-year results from the China PEACE-retrospective CathPCI study. *JAMA Intern Med* 2016;176:512–21.
- 8 Wang C, Li X, Lu FL, *et al.* Comparison of six risk scores for in-hospital mortality in Chinese patients undergoing heart valve surgery. *Heart Lung Circ* 2013;22:612–7.
- 9 Zheng Z, Zhang L, Li X, *et al.* SinoSCORE: a logistically derived additive prediction model for post-coronary artery bypass grafting in-hospital mortality in a Chinese population. *Front Med* 2013;7:477–85.
- 10 Zheng Z, Zhang L, Hu S, *et al.* Risk factors and in-hospital mortality in Chinese patients undergoing coronary artery bypass grafting: analysis of a large multi-institutional Chinese database. *J Thorac Cardiovasc Surg* 2012;144:355–9.
- 11 Li Y, Wang DD, Ley SH, *et al.* Potential impact of time trend of life-style factors on cardiovascular disease burden in China. *J Am Coll Cardiol* 2016;68:818–33.
- 12 Shahian DM, O'Brien SM, Filardo G, *et al.* The Society of Thoracic Surgeons 2008 cardiac surgery risk models: part 1--coronary artery bypass grafting surgery. *Ann Thorac Surg* 2009;88:S2–S22.
- 13 Shahian DM, O'Brien SM, Filardo G, *et al.* The Society of Thoracic Surgeons 2008 cardiac surgery risk models: part 3--valve plus coronary artery bypass grafting surgery. *Ann Thorac Surg* 2009;88:S43–S62.
- 14 Bonaros N, Vill D, Wiedemann D, *et al.* Major risk stratification models do not predict perioperative outcome after coronary artery bypass grafting in patients with previous percutaneous intervention. *Eur J Cardiothorac Surg* 2011;39:e164–e169.
- 15 Pinna-Pintor P, Bobbio M, Colangelo S, *et al.* Inaccuracy of four coronary surgery risk-adjusted models to predict mortality in individual patients. *European Journal of Cardio-Thoracic Surgery* 2002;21:199–204.
- 16 Zhang L, Wang F, Wang L, *et al.* Prevalence of chronic kidney disease in China: a cross-sectional survey. *Lancet* 2012;379:815–22.
- 17 Feng XL, Pang M, Beard J. Health system strengthening and hypertension awareness, treatment and control: data from the China Health and Retirement Longitudinal Study. *Bull World Health Organ* 2014;92:29–41.
- 18 Lu J, Liang L, Feng Y, *et al.* Air pollution exposure and physical activity in china: current knowledge, public health implications, and future research needs. *Int J Environ Res Public Health* 2015;12:14887–97.
- 19 Lim SS, Vos T, Flaxman AD, *et al.* A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012;380:2224–60.
- 20 Yang G, Wang Y, Zeng Y, *et al.* Rapid health transition in China, 1990–2010: findings from the Global Burden of Disease Study 2010. *Lancet* 2013;381:1987–2015.
- 21 Qadir I, Alamzaib SM, Ahmad M, *et al.* EuroSCORE vs. EuroSCORE II vs. Society of Thoracic Surgeons risk algorithm. *Asian Cardiovasc Thorac Ann* 2014;22:165–71.