Validation of a questionnaire measuring patient knowledge of atrial fibrillation in an Asian cohort

Rebecca Hoe,¹ Wanyun Lin,² Mary Ann Cruz Bautista,³ Hubertus Johannes Maria Vrijhoef,^{4,5} Toon Wei Lim⁶

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/heartasia-2018-011143>).

¹Singapore General Hospital, Singapore, Singapore

²National University Hospital, Singapore, Singapore

³Duke-NUS Medical School, Singapore, Singapore

⁴Department of Patient & Care, Maastricht University Medical Center, Maastricht, The Netherlands

⁵Panaxea, Amsterdam, The Netherlands

⁶National University Heart Centre, National University Hospital, Singapore, Singapore

Correspondence to

Dr Toon Wei Lim, National University Heart Centre, National University Hospital, Singapore 119228, Singapore; toon_wei_lim@nuhs.edu.sg

HJMV and TWL contributed equally.

HJMV and TWL are joint senior authors.

Received 6 November 2018

Revised 19 February 2019

Accepted 22 February 2019

ABSTRACT

Background Poor patient understanding of atrial fibrillation (AF) may contribute to underuse of anticoagulation. There are no validated instruments to measure patient knowledge in Asian cohorts. This study aims to validate a disease-specific questionnaire measuring the level of understanding of AF and its treatment among patients with AF in Singapore. **Methods** A 10-item interviewer-administered questionnaire was created based on previously published questionnaires. Face and content validity were assessed. 165 participants were identified by convenience sampling at cardiology clinics of a tertiary hospital. The questionnaire was administered in either English (n = 53) or Mandarin (n = 112). Exploratory factor analysis was performed using principal component method. Internal consistency was evaluated using Cronbach's alpha coefficient.

Results Face validity was tested by surveying 10 cardiologists who could all identify what the questionnaire was designed to measure. Mean content validity ratio across items was 0.9. Participants were 68.7 (SD 10.5) years old. 55.8% were male. 95.2% were on oral anticoagulation. Kaiser-Meyer-Olkin measure was 0.67 and Bartlett's test of sphericity was significant (p < 0.01). Four factors were retained based on the eigenvalue > 1. These were knowledge of the following: disease characteristics, disease-specific treatment, role of treatment in symptom management and treatment mechanisms. Internal consistency was good (Cronbach's alpha = 0.71).

Conclusions A questionnaire on the knowledge of AF and its treatment was validated in a cohort of Asian patients in English and Mandarin. It allows quantification of patient knowledge and may be useful in Asian populations to assess the efficacy of interventions to improve patient understanding of AF.

INTRODUCTION

Atrial fibrillation (AF) is the most common clinically significant heart rhythm disorder in Singapore,¹ with an estimated overall prevalence of 1.5%.² With ageing populations, it is predicted that the number of patients with AF will increase by twofold over the next 50 years.³ This poses a challenge for healthcare systems globally as AF is associated with increased rates of heart failure, cardioembolic stroke⁴ and translates to a poorer quality of life,⁵ greater hospital costs⁶ and higher mortality. There is thus an incentive for healthcare systems to focus on strategies to reduce and prevent the development of complications of AF.

Key messages

What is already known about this subject?

- Patients with atrial fibrillation (AF) have been shown to have poor understanding of their disease and treatment. Objective instruments to measure the level of knowledge of AF among patients have been designed and validated in European populations.

What does this study add?

- This is the first questionnaire that has been validated in English and Mandarin that can be used in an Asian cohort to objectively measure the level of knowledge of AF.

How might this impact clinical practice?

- This knowledge questionnaire can potentially facilitate the development and implementation of patient-centred education interventions, which may lead to improved patient outcomes.

Anticoagulation, be it warfarin or novel oral anti-coagulants (NOACs), is a well-established means of reducing stroke risk.⁷ Warfarin has been shown to reduce stroke risk by 65%, whereas NOACs have been shown to be equal if not more efficacious.⁸⁻⁹ Although the benefits of anticoagulation for stroke prevention in AF are well established, its underutilisation is also widely reported, with less than half of patients with AF receiving anticoagulation.¹⁰ Identifying and overcoming barriers to treatment are areas of active research.

Educational intervention in chronic diseases has been shown to significantly improve knowledge, symptom monitoring, medication use and compliance, thus leading to better disease control. Clark *et al*¹¹ demonstrated in a meta-analysis that the most effective intervention programmes for patients with heart failure involved improving patient knowledge of heart failure, which translated to better disease control. This has been shown in patients with AF as well.¹²⁻¹⁴ Several studies have shown that patients with AF have poor understanding of their disease and its treatment.¹⁵⁻¹⁷ An objective method to measure the level of knowledge among patients with AF would better direct educational resources and help in assessing the efficacy of interventions. Such instruments have previously been studied and validated in European populations.¹⁸

Despite a comparable prevalence of AF in Asian populations and rapidly progressing health literacy, there have been few studies¹⁹ assessing the knowledge of these patients, and no validated instrument

© Author(s) (or their employer(s)) 2019. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Hoe R, Lin W, Bautista MAC, *et al.* *Heart Asia* 2019;**11**:e011143. doi:10.1136/heartasia-2018-011143

that allows us to do so reliably. This study therefore aims to validate an AF knowledge scale, both in Mandarin and in English, for the multiethnic and multilingual Asian population of Singapore.

MATERIALS AND METHODS

Design and development of the questionnaire

The questionnaire was developed in three phases: (1) literature review and first construction, (2) testing for face validity and content validity and (3) factor analysis and testing of internal consistency. The scope of this work is limited to face and content validity testing as well as construct validity testing.

The questionnaire was created after a literature review resulting in in-depth analysis of four previous studies by Hendriks *et al.*,²⁰ Aliot *et al.*,²¹ Koponen *et al.*²² and Lane *et al.*,¹² which all aimed to assess knowledge of AF in patients.

The questionnaire was designed to encompass the following: (1) knowledge of the disease and (2) knowledge of treatment. The questionnaire (figure 1) consists of 10 items that were grouped into the following categories: AF in general (two items: Q1 and Q2), symptoms of AF (one item: Q3) and AF treatment including aims, options and complications (seven items: Q4–Q10). A Mandarin version (online supplementary file) was developed through forward and backward translation with accredited bilingual translators working independently.

Patient demographic data were collected by interviewing patients and through electronic medical records. These included age, gender, education level and medical history pertaining to duration of AF, duration of anticoagulation, frequency of follow-up, number of hospitalisations due to AF and complications of AF, most recent international normalised ratio, and presence of other medical comorbidities (constituents of the CHA2DS2-VASc score).

Validity

Face and content validity were assessed by presenting the questionnaire to 10 cardiologists from a single tertiary hospital who were active in the clinical care of patients with AF.

Face validity was tested by surveying the cardiologists on the apparent intent of the questionnaire while content validity was measured using Lawshe's Content Validity Ratio²³ by inviting them to label each item of the questionnaire as 'essential', 'useful but not essential' or 'not useful or essential'. Each of them was also asked for their opinion regarding the readability and comprehensibility of the questionnaire as a whole. The items of the questionnaire were then reworded based on the feedback obtained.

Construct validity was determined by performing exploratory factor analysis with varimax rotation. For the set of 10 items contained in the questionnaire, two measures were obtained to determine the suitability of performing a factor analysis: the value of the Kaiser-Meyer-Olkin (KMO) measure and the statistical significance of the Bartlett's test of sphericity. Obtaining an adequate KMO value (ie, >0.5) and rejecting the null hypothesis (ie, statistically significant Bartlett's test of sphericity value) would indicate that performing a factor analysis was appropriate. The KMO test measures the sampling adequacy and is a measure of the strength of the relationship between variables. The KMO measure should be greater than 0.5 for factor analysis to proceed. The null hypothesis is that the correlation matrix is an identity matrix. The null hypothesis also needs to be rejected before proceeding with factor analysis. Rejection of the null hypothesis suggests that all the variables correlate sufficiently to provide an adequate basis for factor analysis. After determining

the suitability for factor analysis, the factor analysis by principal component method with varimax rotation (which assumes no correlation between the factors in the analysis) was completed and factor loadings of at least 0.3 were considered acceptable.²⁴

Reliability

Internal consistency reliability was evaluated using Cronbach's alpha coefficient. An acceptable value for this coefficient is at least 0.70. A value between 0.15 and 0.50 confirms the unidimensionality of the scale.

Participant recruitment

A total of 165 participants with AF were identified by convenience sampling in the cardiology outpatient clinics of a tertiary academic hospital. All patients with a known diagnosis of AF were approached to participate in the questionnaire. Of these, 53 participants completed the questionnaire in English and 112 completed it in Mandarin based on their preference of language. The questionnaire was verbally administered by trained interviewers who had undergone a standardisation course.

Statistics

Statistical analysis was performed using SPSS (V.22, IBM Corporation, Armonk, NY, USA). Demographic characteristics were recorded and divided into categorical variables. Descriptive statistics were used to present the frequencies of demographic characteristics within each category, which were subsequently expressed as a percentage of the total number of respondents per characteristic. Similarly, responses for items 1–9 in the questionnaire were dichotomised into 'correct' and 'mistaken/do not know' categories. These were analysed using descriptive statistics as well.

Statement of ethical publishing

Ethics approval for this study was obtained from the National Healthcare Group, Domain Specific Review Board, Singapore, prior to participant recruitment and data collection and analysis. Written consent was obtained from all participants.

RESULTS

Face and content validity

All 10 cardiologists surveyed identified correctly that the questionnaire intended to measure the level of understanding of AF and its treatment. Mean content validity ratio across items according to Lawshe's content validity rule was 0.9 which was consistent with good content validity.

Demographics

Interview participants were 68.7 (SD 10.5) years old with a slight preponderance of men (table 1). Two-thirds of them did not progress beyond primary school education. Median CHA2DS2-VASc score was 4 (25th–75th centiles 3–5), 92.0% who had a score of 2 and above. There was a high prevalence of hypertension and diabetes. Most (95.2%) were on oral anticoagulation. All 165 patients were able to complete the interview.

Construct validity

Exploratory factor analysis by principal component method yielded a KMO measure of 0.67. The test of hypothesis using the Bartlett's test of sphericity was significant (p value < 0.01). Four factors were retained based on Kaiser's criterion that suggests retention of factors with eigenvalues greater than 1.²⁵ Altogether, these four factors explained 64.1% of the variance.

ATRIAL FIBRILLATION KNOWLEDGE, ATTITUDE AND PERCEPTIONS QUESTIONNAIRE

National University of Singapore and National University Health System

NAME OF INVESTIGATOR:		DATE AND TIME:		
PATIENT DEMOGRAPHICS				
PATIENT SERIAL NUMBER: DOB: AGE: GENDER: M/F		EDUCATION LEVEL: - LESS THAN PRIMARY SCHOOL ☐ A LEVELS ☐ - PRIMARY SCHOOL ☐ UNDERGRADUATE ☐ - O LEVELS ☐ POSTGRADUATE ☐ - POLYTECHNIC DEGREE ☐		
MEDICAL HISTORY				
ATRIAL FIBRILLATION (AF) HISTORY DURATION OF AF: DURATION OF ANTICOAGULATION: FREQUENCY OF FOLLOWUP AT NUH: NUMBER OF HOSPITALISATIONS DUE TO AF/COMPLICATIONS OF AF: MOST RECENT INR:		OTHER MEDICAL COMORBIDITIES: - HYPERTENSION ☐ - DIABETES MELLITUS ☐ - CONGESTIVE HEART FAILURE/ LEFT VENTRICULAR SYSTOLIC DYSFUNCTION ☐ - STROKE ☐ - TIA ☐ - THROMBOEMBOLISM ☐ - PERIPHERAL ARTERIAL DISEASE ☐ - MYOCARDIAL INFARCTION ☐ - ATHEROSCLEROSIS ☐		CHA ₂ DS ₂ VASc SCORE: C – H – A ₂ – D – S ₂ – V – A – Sc –
KNOWLEDGE				
OF ATRIAL FIBRILLATION				
1	WHAT IS THE NAME OF YOUR HEART CONDITION?	C	DK	MISTAKEN:
	IF PATIENT SAYS NAME OF ANOTHER HEART CONDITION, ASK "ANY OTHER HEART CONDITION?"	C	DK	MISTAKEN:
2	WHAT IS ATRIAL FIBRILLATION?	C	DK	MISTAKEN
3	IS THIS A SYMPTOM OF ATRIAL FIBRILLATION? - IRREGULAR HEART BEAT - BREATHLESSNESS - CHEST DISCOMFORT - LIGHT-HEADEDNESS - DECREASED EXERCISE ABILITY	C C C C C	DK DK DK DK DK	MISTAKEN MISTAKEN MISTAKEN MISTAKEN MISTAKEN
4	WHAT ARE WE TRYING TO PREVENT WHEN WE TREAT ATRIAL FIBRILLATION? - SYMPTOMATIC RELIEF - STROKE - HEART FAILURE - HEART ATTACK	C C C C	DK DK DK DK	MISTAKEN MISTAKEN MISTAKEN MISTAKEN
OF TREATMENT				
5	ARE YOU TAKING ANY BLOOD THINNING MEDICATION?	C	DK	MISTAKEN
6	WHAT IS THE NAME OF YOUR BLOOD THINNING MEDICATION?	C	DK	MISTAKEN
7	WHY ARE YOU ON A BLOOD THINNING MEDICATION?	C	DK	MISTAKEN
8	IS THIS A COMPLICATION OF BLOOD THINNING MEDICATION? - BRUISING - BLEEDING	C C	DK DK	MISTAKEN MISTAKEN
9	DO YOU KNOW HOW YOUR BLOOD IS BEING MONITORED?	C	DK	MISTAKEN
10	OTHER THAN BLOOD THINNING MEDICATION, WHAT OTHER TREATMENTS ARE THERE FOR ATRIAL FIBRILLATION? (OEQ) - RATE/RHYTHM CONTROL DRUGS - RADIOFREQUENCY ABLATION - CARDIOVERSION - OTHERS: _____	C C C C		To prompt 1x by asking "any others?" if participant does not spontaneously offer all NOT OFFERED NOT OFFERED NOT OFFERED NOT OFFERED

Figure 1 English version of questionnaire. AF, atrial fibrillation; INR, international normalised ratio; NUH, National University Hospital; TIA, transient ischaemic attack.

Table 1 Demographic characteristics of total patient population

Demographic	Number	Percentage (%)
Age (years)		
<65	50	30.3
65–74	62	37.6
75 and above	53	32.1
Gender		
Male	92	55.8
Female	73	44.2
Educational level		
Less than primary school	67	40.6
Completed primary school	43	26.1
Completed O Levels	31	18.8
Completed A Levels	4	2.4
Completed polytechnic degree	9	5.5
Completed undergraduate	8	4.8
Completed postgraduate	1	0.6
Duration of medication use		
Total number on medication	157	95.2
<1 year	55	33.3
1–5 years	62	37.6
5–10 years	25	15.2
>10 years	15	9.1
Risk factors for stroke		
Hypertension	129	78.2
Diabetes mellitus	66	40.0
Congestive heart failure	59	35.8
Previous stroke	26	15.8
Previous TIA	15	9.1
Thromboembolism	0	0.0
Peripheral arterial disease	25	15.2
Acute coronary syndrome	36	21.8
Hyperlipidaemia	111	67.3
Median CHA2DS2VAsC score	4	

TIA, transient ischaemic attack.

Factors 1–4 explained 29.3%, 12.7%, 11.9% and 10.3% of the variance, respectively.

The rotated component matrix (table 2) shows the factor loadings for each component (factor), taking note of all factor

Table 2 Rotated component matrix

	Component/factor			
	1	2	3	4
Item 1	0.026	0.171	0.774	–0.065
Item 2	0.182	–0.058	0.831	0.013
Item 3	–0.149	0.490	0.222	0.219
Item 4	0.041	0.110	–0.015	0.890
Item 5	0.158	0.735	–0.160	–0.011
Item 6	0.203	0.600	0.148	–0.313
Item 7	0.307	0.627	0.097	0.118
Item 8	0.387	0.323	0.087	–0.367
Item 9	0.951	0.135	0.064	–0.025
Item 10	0.944	0.172	0.144	0.002

Extraction method: factor analysis by component method.

Rotation method: varimax with Kaiser normalisation.

Factor loadings higher than 0.3 are in bold.

* Rotation converged in six iterations.

loadings higher than 0.3. The higher the absolute value of the loading, the greater that item contributes to the factor. Three items (Q8, Q9 and Q10) loaded on Factor 1; four items (Q3, Q5, Q6 and Q7) on Factor 2; two items (Q1 and Q2) on Factor 3 and one item (Q4) on factor 4. These four factors were named as follows: Factor 1 (Q8, Q9 and Q10), knowledge of treatment mechanisms; Factor 2 (Q3, Q5, Q6 and Q7), knowledge of disease-specific treatment; Factor 3 (Q1 and Q2), knowledge of disease characteristics and Factor 4 (Q4), role of treatment in symptom management.

Reliability

Internal consistency was good (Cronbach's $\alpha=0.71$), indicating that all items were correlated. Removal of item 4 improved the Cronbach's α to 0.73; nevertheless, this item was retained in view of its value to daily practice and the small difference in the value of Cronbach's α .

Questionnaire responses

Responses are detailed in table 3.

Knowledge of disease characteristics

Participants generally demonstrated poor knowledge of the disease, with a minority (16.4%) aware of the name of the heart condition they had and were on follow-up for. When offered the term 'atrial fibrillation', few (16.4%) knew what it was. However, majority of participants were able to identify common symptoms of their cardiac condition.

Knowledge of role of treatment in symptom management

Majority of participants identified that treatment is directed towards cardiac complications of AF (symptomatic relief and heart failure). However, they were also of the mistaken belief that treatment of AF lowers risk of myocardial infarction. Notably, only 12.7% were aware that one of the main aims of AF treatment was stroke prevention.

Knowledge of disease-specific treatment

Most participants (92.7%) correctly stated whether or not they were on anticoagulation. Of these, only 54.5% named their anti-coagulant correctly.

Knowledge of treatment mechanisms

A majority of participants were able to correctly identify complications of anticoagulation and that the adequacy of anticoagulation was being monitored by blood tests. There was overall poor understanding of alternative therapies available for AF, with 75.3% not knowing of any other forms of therapy apart from anticoagulation.

DISCUSSION

This paper reports the development and validation of a questionnaire that can be used in an Asian cohort to objectively measure the level of knowledge of AF. Importantly, this questionnaire was administered in English and Mandarin, allowing the objective measurement of the level of knowledge among patients with AF in a multilingual, multiethnic Asian population. The participants' responses also highlight the low levels of knowledge among patients with AF which may represent a potential target for intervention.

The questionnaire had a high mean content validity ratio. The four factors identified using factor analysis appear to be subsets of the original two components of knowledge the questionnaire

Table 3 Participant responses

Item	Question asked	Patient answer	Percentage (%)
1	What is the name of your heart condition?	Correct	16.4%
		Mistaken	40.6
		Do not know/no answer provided	43.0
2	What is atrial fibrillation?	Correct	16.4
		Mistaken/do not know	82.4
3	Is this a symptom of atrial fibrillation?	Correct	68.5
		Mistaken/do not know	31.5
	Irregular heart beat	Correct	60.3
		Mistaken/do not know	39.5
	Breathlessness	Correct	52.1
		Mistaken/do not know	47.9
	Chest discomfort	Correct	42.4
		Mistaken/do not know	57.6
4	Light-headedness	Correct	63.6
		Mistaken/do not know	36.4
	Decreased exercise ability	Correct	63.0
		Mistaken/do not know	37.0
	What are we trying to prevent when we treat atrial fibrillation?	Correct	12.7
		Mistaken/do not know	87.3
	Symptomatic relief	Correct	57.6
		Mistaken/do not know	42.4
5	Stroke	Correct	43.6
		Mistaken/do not know	56.4
6	Heart failure	Correct	92.7
		Mistaken/do not know	7.3
7	Heart attack	Correct	54.5
		Mistaken/do not know	45.5
8	Are you taking any blood thinning medication?	Correct	68.5
		Mistaken/do not know	31.5
9	What is the name of your blood thinning medication?	Correct	68.5
		Mistaken/do not know	31.5
	Why are you on a blood thinning medication?	Correct	68.5
		Mistaken/do not know	31.5
10	Is this a complication of blood thinning medication?	Correct	68.5
		Mistaken/do not know	31.5
	Bruising	Correct	74.5
		Mistaken/do not know	25.5
11	Bleeding	Correct	66.1
		Mistaken/do not know	33.9
12	Do you know how your blood is being monitored?	Correct	11.5
		Mistaken/do not know	7.3
	Other than blood thinning medications, what other treatments are there for atrial fibrillation?	Radiofrequency ablation	1.2
		Cardioversion	11.5

was designed to assess. Internal consistency of the questionnaire was good in spite of a small number of items. The results are comparable to the validated AF knowledge scale developed by Hendriks *et al* which has a Cronbach's alpha of 0.58 and good face, content and construct validity.²⁰

Thus far, this is the only study that has validated an instrument to assess patient's knowledge of AF in a multilingual Asian patient population. It is the first AF knowledge questionnaire validated in Mandarin. As this is a short questionnaire of 10 items, it can be administered by trained clinic staff within 5 min prior to consultations.

The knowledge this cohort of patients with AF had of their disease, its complications and its treatment was generally poor. Similar findings suggesting poor patient knowledge of AF have also been shown by other studies. Lip *et al*¹⁸ showed that only 63% of 119 patients studied were aware of their cardiac condition and only 52% were aware of the reasons for initiating oral anticoagulation therapy. The AF aware group also showed in 2010 that 25% of 825 patients with AF studied were unable to explain AF.²¹ Due to different methods of testing for patient knowledge, direct comparison of results with these studies is not possible, but it highlights that this study's findings are not unique and probably typical of patients with AF worldwide.

The low level of patient knowledge of AF raises several possible gaps, such as physician communication with patients, language barriers (especially in a multiethnic society) and the failure to allocate adequate time and resources to patient education. In this study, the relatively low level of formal education may also be a contributing factor. It does, however, point to important targets for therapeutic interventions for improving patient knowledge to promote patient empowerment and enable self-management, which may, in turn, translate into improved compliance with treatment.¹² Studies by Howitt *et al*¹³ and Protheroe *et al*¹⁴ showed that patient's knowledge and beliefs about their disease and treatment are important factors influencing their compliance with anticoagulation. As such, various centres managing AF have invested in specialised AF clinics which dedicate resources to patient education to promote 'self-management' of AF.^{26 27}

An instrument that can measure patients' level of knowledge of a disease is important to identify patients with the greatest education need and evaluate the effect of education interventions. Previously validated questionnaires by Koponen *et al*²² and Hendriks *et al*²⁰ were written and administered in English and Dutch, respectively to a Western population and may not be fully applicable to the local population in Singapore where a large proportion of older patients are unable to communicate in English.

This study has several limitations. First, with regard to questionnaire administration, an interviewer is required to administer the questionnaire, necessitating additional manpower and potentially resulting in differences in interpretation of patient responses. This was minimised by ensuring interviewers undergo a standardisation course. The questionnaire was also designed in a manner that requires the interviewer to review the participants' medical records to determine whether their responses are correct, requiring more effort on the part of the interviewer. Second, while English and Mandarin are the most commonly spoken languages, there is a significant proportion of patients in the population that do not speak either. Third, the questionnaire was only tested in a single centre in an outpatient setting. When using the questionnaire in a different context, it is advised to further validate the questionnaire. For example, its face validity may be further increased by including different experts. It would also be useful to test the relationship between disease knowledge and other variables (eg, self-management behaviour) to assess the usefulness of the questionnaire to obtain treatment objectives. Finally, the validation process was limited to face and content validity testing as well as construct validity testing. Confirmatory factor analysis is an additional step that can be performed. In addition, test-retest reliability and agreement statistics would be important psychometric properties to establish. The methodological quality of the questionnaire can also be further assessed in accordance with the nsensus-based standards for the selection of health status measurement instruments checklist.²⁸

CONCLUSIONS

The present study reports the development and validation of a questionnaire for measuring patient knowledge of AF in Singapore. As this is the first AF knowledge questionnaire validated in Mandarin, it is of relevance to the Asia Pacific region, both in predominantly Mandarin-speaking countries and also in countries with significant numbers of Chinese immigrants. This knowledge questionnaire can potentially facilitate the development and implementation of patient-centred education interventions, which may lead to improved patient-related outcomes such as patient empowerment and self-management. The low levels of patient knowledge found in this study suggest the pressing need to improve this aspect of AF management.

Acknowledgements We would like to thank the patients, nurses and cardiologists for their participation and support. A special word of thanks to Dr Leong Wai Siang for his contribution to the development of the Mandarin questionnaire.

Contributors All authors were involved in the conception and design of the work, data analysis and interpretation, critical revision of the manuscript, and final approval of the version to be published. WL and RH were responsible for data collection and drafting the manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- Ministry of Health. *Management of atrial fibrillation. Clinical practice guidelines*. Singapore: Ministry of Health, 2004.
- Yap KB, Ng TP, Ong HY. Low prevalence of atrial fibrillation in community-dwelling Chinese aged 55 years or older in Singapore: a population-based study. *J Electrocardiol* 2008;41:94–8.
- Go AS, Hylek EM, Phillips KA, et al. Prevalence of diagnosed atrial fibrillation in adults: national implications for rhythm management and stroke prevention: the anticoagulation and risk factors in atrial fibrillation (atria) study. *JAMA* 2001;285:2370–5.
- Stewart S, Hart CL, Hole DJ, et al. A population-based study of the long-term risks associated with atrial fibrillation: 20-year follow-up of the Renfrew/Paisley study. *Am J Med* 2002;113:359–64.
- Thrall G, Lip GYH, Carroll D, et al. Depression, anxiety, and quality of life in patients with atrial fibrillation. *Chest* 2007;132:1259–64.
- Ringborg A, Nieuwlaar R, Lindgren P, et al. Costs of atrial fibrillation in five European countries: results from the Euro Heart survey on atrial fibrillation. *Europace* 2008;10:403–11.
- Hart RG, Benavente O, McBride R, et al. Antithrombotic therapy to prevent stroke in patients with atrial fibrillation: a meta-analysis. *Ann Intern Med* 1999;131:492–501.
- Madzak A, Larsen TB, Lane DA, et al. Composite end point analyses of non-vitamin K antagonist oral anticoagulants compared with Warfarin in patients with atrial fibrillation. *Expert Rev Cardiovasc Ther* 2015;13:1155–63.
- Dentali F, Riva N, Crowther M, et al. Efficacy and safety of the novel oral anticoagulants in atrial fibrillation: a systematic review and meta-analysis of the literature. *Circulation* 2012;126:2381–91.
- Bungard TJ, Ghali WA, Teo KK, et al. Why do patients with atrial fibrillation not receive warfarin? *Arch Intern Med* 2000;160:41–6.
- Clark AM, Wiens KS, Banner D, et al. A systematic review of the main mechanisms of heart failure disease management interventions. *Heart* 2016;102:707–11.
- Lane DA, Ponsford J, Shelley A, et al. Patient knowledge and perceptions of atrial fibrillation and anticoagulant therapy: effects of an educational intervention programme. the West Birmingham atrial fibrillation project. *Int J Cardiol* 2006;110:354–8.
- Howitt A, Armstrong D. Implementing evidence based medicine in general practice: audit and qualitative study of antithrombotic treatment for atrial fibrillation. *BMJ* 1999;318:1324–7.
- Protheroe J, Fahey T, Montgomery AA, et al. The impact of patients' preferences on the treatment of atrial fibrillation: observational study of patient based decision analysis. *BMJ* 2000;320:1380–4.
- Desteghe L, Engelhard L, Raymaekers Z, et al. Knowledge gaps in patients with atrial fibrillation revealed by a new validated knowledge questionnaire. *Int J Cardiol* 2016;223:906–14.
- Alphonsa A, Sharma KK, Sharma G, et al. Knowledge regarding oral anticoagulation therapy among patients with stroke and those at high risk of thromboembolic events. *J Stroke Cerebrovasc Dis* 2015;24:668–72.
- Smith MB, Christensen N, Wang S, et al. Warfarin knowledge in patients with atrial fibrillation: implications for safety, efficacy, and education strategies. *Cardiology* 2010;116:61–9.
- Lip GYH, Kamath S, Jafri M, et al. Ethnic differences in patient perceptions of atrial fibrillation and anticoagulation therapy: the West Birmingham atrial fibrillation project. *Stroke* 2002;33:238–42.
- Lee VVY, Tam CS, Yan BP, et al. Barriers to warfarin use for stroke prevention in patients with atrial fibrillation in Hong Kong. *Clin Cardiol* 2013;36:166–71.
- Hendriks JML, Crijns HJGM, Tieleman RG, et al. The atrial fibrillation knowledge scale: development, validation and results. *Int J Cardiol* 2013;168:1422–8.
- Aliot E, Breithardt G, Brugada J, et al. An international survey of physician and patient understanding, perception, and attitudes to atrial fibrillation and its contribution to cardiovascular disease morbidity and mortality. *Europace* 2010;12:626–33.
- Koponen L, Rekola L, Ruotsalainen T, et al. Patient knowledge of atrial fibrillation: 3-month follow-up after an emergency room visit. *J Adv Nurs* 2008;61:51–61.
- Lawshe CH. A quantitative approach to content validity. *Pers Psychol* 1975;28:563–75.
- Brown JD. Questions and answers about language testing statistics: choosing the right number of components or factors in pCa and EFA. *Shiken: JALT Testing & Evaluation SIG Newsletter* 2009;13:19–23.
- Ledesma RD, Valero-Mora P. Determining the number of factors to retain in EFA: an easy-to-use computer program for carrying out Parallel analysis. *Pract Assess Res Eval* 2007;12:1–11.
- Tran HN, Tafreshi J, Hernandez EA, et al. A multidisciplinary atrial fibrillation clinic. *Curr Cardiol Rev* 2013;9:55–62.
- Qvist I, Hendriks JML, Møller DS, et al. Effectiveness of structured, hospital-based, nurse-led atrial fibrillation clinics: a comparison between a real-world population and a clinical trial population. *Open Heart* 2016;3:e000335.
- Mokkink LB, Terwee CB, Patrick DL, et al. The COSMIN checklist for assessing the methodological quality of studies on measurement properties of health status measurement instruments: an international Delphi study. *Qual Life Res* 2010;19:539–49.