spectroscopy, or NIRS, in the monitoring of lower limb perfusion during ECMO therapies.

Strategies alternative to DPC include end-to-side graft, posterior tibial artery retrograde perfusion, axillary cannulation and central sports mode. Novel bidirectional perfusion cannula may appear as a future promising option.

REFERENCES

- Cheng R, Hachamovitch R, Kittleson M, Patel J, Arabia F, Moriguchi J, Esmailian F, Azarbal B. Complications of extracorporeal membrane oxygenation for treatment of cardiogenic shock and cardiac arrest: a meta-analysis of 1,866 adult patients. *Ann Thorac Surg* 2014;**97**:610–616.
- Tanaka D, Hirose H, Cavarocchi N, Entwistle JW. The impact of vascular complications on survival of patients on venoarterial extracorporeal membrane oxygenation. *Ann Thorac Surg* 2016;**101**:1729–1734.
- Juo YY, Skancke M, Sanaiha Y, Mantha A, Jimenez JC, Benharash P. Efficacy of distal perfusion cannulae in preventing limb ischemia during extracorporeal membrane oxygenation: a systematic review and meta-analysis. *Artif Organs* 2017;**41**:E263–E273.
- Wong JK, Smith TN, Pitcher HT, Hirose H, Cavarocchi NC. Cerebral and lower limb near-infrared spectroscopy in adults on extracorporeal membrane oxygenation. *Artif Organs* 2012;**36**:659–667.
- Spurlock DJ, Toomasian JM, Romano MA, Cooley E, Bartlett RH, Haft JW. A simple technique to prevent limb ischemia during veno-arterial ECMO using the femoral artery: the posterior tibial approach. *Perfusion* 2012;**27**:141–145.
- Marasco SF, Tutungi E, Vallance SA, Udy AA, Negri JC, Zimmet AD, McGiffin DC, Pellegrino VA, Moshinsky RA. A Phase 1 study of a novel bidirectional perfusion cannula in patients undergoing femoral cannulation for cardiac surgery. *Innovations (Phila)* 2018;**13**:97–103.

28 SHOULD WE BE USING DIGOXIN IN 2018?

Ganesan Karthikeyan. All India Institute of Medical Sciences, New Dehli, India

10.1136/heartasia-2019-apahff.28

Digoxin is often used in the management of patients with heart failure (HF) with or without atrial fibrillation (AF). There is sound biological rationale for the use of digoxin, but the data on clinical outcomes with digoxin use in this patient population are conflicting. There is a single adequately powered randomised trial of digoxin in patients with HF in sinus rhythm.¹ This trial showed a small but significant reduction in the rate of hospitalisation due to HF with the use of digoxin, when compared to placebo, among patients treated with ACE inhibitors and diuretics. There was no effect on mortality.¹ On the contrary, there are numerous secondary analyses of observational data from randomised trials which suggest that there may be an increased risk of death from using digoxin, both in patients with HF, those in AF, or both.² However, observational data on digoxin use suffer from treatment bias (confounding by indication), as the sickest patients are the ones who are prescribed digoxin. Propensity matched analyses have been attempted to overcome the effect of this bias with conflicting results.^{2–3} However, it is likely that because the magnitude of this bias is large, no amount of statistical adjustment can yield reliable effect estimates. This highlights the need for large, randomised trials of digoxin.³

REFERENCES

- Digitalis Investigation G. The effect of digoxin on mortality and morbidity in patients with heart failure. *N Engl J Med* 1997;**336**:525–533.
- Ziff OJ, Lane DA, Samra M, Griffith M, Kirchhof P, Lip GY, Steeds RP, Townend J, Kotecha D. Safety and efficacy of digoxin: systematic review and meta-analysis of observational and controlled trial data. *BMJ* 2015;**351**:h4451.
- Karthikeyan G, Devasenapathy N, Zühlke L, Engel ME, Rangarajan S, Teo KK, Mayosi BM, Yusuf S; Global Rheumatic Heart Disease Registry (REMEDI) Investigators. Digoxin and clinical outcomes in the Global Rheumatic Heart Disease Registry. *Heart* 2018 Sep 12. pii: heartjnl-2018-313614.

29 MANAGEMENT OF PULMONARY HYPERTENSION IN SYSTEMIC LUPUS ERYTHEMATOSUS PATIENTS

Priscilla CH Wong. Department of Medicine and Therapeutics, Prince of Wales Hospital, Hong Kong SAR

10.1136/heartasia-2019-apahff.29

Pulmonary arterial hypertension (PAH) is a complex and devastating disease. According to a longitudinal United States-based registry, connective tissue disease (CTD) accounted for more than 50% of all patients with PAH, in which systemic sclerosis comprised the largest CTD-related PAH.¹ In contrast to Western countries, systemic lupus erythematosus (SLE) is a more common CTD than systemic sclerosis in the Asia-Pacific region.² A cohort study from China has shown that SLE, instead of systemic sclerosis, comprised the largest proportion of all CTD-related PAH.³ The prevalence of PAH in SLE is estimated at 0.5%–17.5%.⁴ The pathogenesis of PAH involves multiple mechanisms including vasculitis, *in situ* thrombosis to interstitial lung disease which may all increase pulmonary vascular resistance and lead to right heart failure. The leading risk factors for the development of PAH in SLE patients include Raynaud's phenomenon, anti-U1 RNP antibody and anti-cardiolipin antibodies positivity. Since PAH is potentially life-threatening, early detection is crucial to improve the outcomes of this condition. Currently, the diagnostic algorithm for PAH in SLE patients follows that of international guidelines. Diagnosis is confirmed by right heart catheterisation. Treatments are similar to the therapeutic interventions for patients with idiopathic PAH. Since inflammatory and dysregulated immune components may play a major role in the pathogenesis of PAH in SLE, glucocorticoids and immunosuppressive therapies including cyclophosphamide are used, although the immunosuppressive therapy trials were small, uncontrolled studies only. Regular follow-up with prognostic evaluation and risk assessment should be performed and the treatment should be individualised accordingly.

REFERENCES

- McGoon MD, Miller DP. REVEAL: a contemporary US pulmonary arterial hypertension registry. *Eur Respir Rev* 2012;**21**:8–18.
- Jakes RW, Bae SC, Louthrenoo W, Mok CC, Navarra SV, Kwon N. Systematic review of the epidemiology of systemic lupus erythematosus in the Asia-Pacific region: prevalence, incidence, clinical features, and mortality. *Arthritis Care Res (Hoboken)* 2012;**64**:159–168.
- Zhao J, Wang Q, Liu Y, Tian Z, Guo X, Wang H, Lai J, Huang C, Yang X, Li M, Zeng X. Clinical characteristics and survival of pulmonary arterial hypertension associated with three major connective tissue diseases: A cohort study in China. *Int J Cardiol* 2017;**236**:432–437.
- Condliffe R, Kiely DG, Peacock AJ, Corris PA, Gibbs JS, Vrapai F, Das C, Elliot CA, Johnson M, DeSoyza J, Torpy C, Goldsmith K, Hodgkins D, Hughes RJ, Pepke-Zaba J, Coghlan JG. Connective tissue disease-associated pulmonary arterial hypertension in the modern treatment era. *Am J Respir Crit Care Med* 2009;**179**:151–157.

30 ADVANCES IN MEDICAL AND INTERVENTIONAL TREATMENTS FOR CTEPH

Yu Taniguchi. Department of Cardiology and Cardiovascular Intervention Unit, Kobe University Hospital, Kobe, Japan

10.1136/heartasia-2019-apahff.30

Pulmonary endarterectomy is the standard care for patients with chronic thromboembolic pulmonary hypertension (CTEPH), however, about 40% of them are inoperable.